Environmental Protection Agency

§ 180.522 Fumigants for processed grains used in production of fermented malt beverage; tolerances for residues.

- (a) *General*. Fumigants for processed grain may be safely used, in accordance with the following conditions.
- (1) *Methyl bromide*. Total residues of inorganic bromides (calculated as Br) from the use of this fumigant shall not exceed 125 parts per milion.
- (2) Methyl bromide is used to fumigate corn grits and cracked rice in the production of fermented malt beverage.
- (3) To assure safe use of the fumigant, its label and labeling shall conform to the label and labeling registered by the U.S. Environmental Protection Agency, and the usage employed should conform with such label or labeling.
- (4) The total residue of inorganic bromides in fermented malt beverage, resulting from the use of corn grits and cracked rice fumigated with the fumigant described in paragraph (a)(2) of this section plus additional residues of inorganic bromides that may be present from uses in accordance with other regulations in this chapter promulgated under section 408 and/or 409 of the Act, does not exceed 25 parts per million bromide (calculated as Br).
- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[71 FR 74818, Dec. 13, 2006]

§ 180.523 Metaldehyde; tolerances for residues.

(a) General. Tolerances are established for residues of the molluscicide metaldehyde in or on food commodities, as follows:

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[73 FR 54963, Sept. 24, 2008]

§ 180.525 Resmethrin; tolerances for residues. (a) General. Tolerances are estab-

lished for residues of the insecticide resmethrin [5-(phenylmethyl)-3methyl 2,2-dimethyl-3-(2furanv11 methyl-1-propenyl) cyclopropanecarboxylate in or on food commodities at 3.0 ppm resulting from use of the insecticide in food handling and storage areas as a space concentration for spot/or crack and crevice treatment and shall be limited to a maximum of 3.00 percent of the active ingredient by weight, and as a space treatment shall be limited to a maximum of 0.5 fluid ounce of 3.0 percent active ingredient by weight per 1000 cubic feet of space provided that the food is removed or covered prior to such use. To assure safe use of the additive, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency, and shall be used in accordance with such label and labeling.

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[71 FR 74819, Dec. 13, 2006]

§ 180.526 Synthetic isoparaffinic petroleum hydrocarbons; tolerances for residues.

(a) General. Synthetic isoparaffinic petroleum hydrocarbons complying with 21 CFR 172.882 (a) and (b) may be safely used as a component of insecticide formulations for use on animal feed in an amount no greater than reasonably required to accomplish its intended effect as an adjuvant in the insecticide formulation and shall not be intended to accomplish any effect in animal feed. It is used or intended for

§ 180.527

use as a component of insecticide formulations used in compliance with regulations issued in 40 CFR part 180 and in this part.

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[40 FR 14161, Mar. 28, 1975, as amended at 50 FR 2959, Jan. 23, 1985, and amended at 53 FR 24668, 24669, June 29, 1988. Redesignated and amended at 63 FR 34319, June 24, 1998]

§ 180.527 Flufenacet, N-(4-fluorophenyl)-N-(1-methylethyl)-2-[[5-(trifluoromethyl)-1, 3, 4-thiadiazol-2-yl] oxylacetamide and its metabolites containing the 4-fluoro-N-methylethyl benzenamine tolerances for residues.

(a) General. Tolerances are established for the combined residues of the herbicide flufenacet, N-(4-fluorophenyl)-N-(1-methylethyl)-2-[[5-(trifluoromethyl)-1, 3, 4-thiadiazol-2-yl] oxy]acetamide and its metabolites containing the 4-fluoro-N-methylethyl benzenamine moiety in or on the following commodities.

Commodity	Parts per million
Cattle, kidney	0.05
Corn, field, forage	0.4
Corn, field, grain	0.05
Corn, field, stover	0.4
Corn, sweet, forage	0.45
Corn, sweet, kernel plus cob with husks re-	
moved	0.05
Corn, sweet, stover	0.30
Goat, kidney	0.05
Hog, kidney	0.05
Horse, kidney	0.05
Sheep, kidney	0.05
Soybean, seed	0.1
Wheat, bran	0.80
Wheat, forage	6.0
Wheat, grain	0.60
Wheat, hay	1.2
Wheat, straw	0.35

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. Tolerances are established for combined residues of flufenacet, N-(4-fluorophenyl)-N-(1-methylethyl)-2-[[5-(trifluoromethyl)-1, 3, 4-thiadiazol-2-yl] oxy]acetamide, and its metabolites containing the 4-fluoro-N-methylethyl benzenamine moiety, with regional registration

Commodity	Parts per million
Grass, forage	7.0 0.4

(d) Indirect or inadvertent residues. Tolerances are established for indirect or inadvertent residues of the herbicide flufenacet, N-(4-fluorophenyl)-N-(1-methylethyl)-2-[[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]oxy]acetamide and its metabolites containing the 4-fluoro-N-methylethyl benzenamine moiety in or on the following raw agricultural commodities when present therein as a result of application of flufenacet to the growing crops in paragraph (a) of this section.

Commodity	Parts per million
Alfalfa, forage	0.1
Alfalfa, hay	0.1
Alfalfa, seed	0.1
Clover, forage	0.1
Clover, hay	0.1
Grain, cereal, group 15, except rice	0.1
16, except rice	0.1
Grass, forage, fodder, and hay, group 17	0.1

[63 FR 26473, May 13, 1998, as amended by 63 FR 50791, Sept. 23, 1998; 64 FR 42846, Aug. 6, 1999; 65 FR 64366, Oct. 27, 2000; 68 FR 2247, Jan. 16, 2003; 68 FR 37759, June 25, 2003; 70 FR 37696, June 30, 2005; 71 FR 76200, Dec. 20, 2006; 72 FR 26310, May 9, 2007]

§ 180.530 2,2-Dimethyl-1,3-benzodioxol-4-ol methylcarbamate; tolerances for residues.

- (a) General. (1) The insecticide 2,2-dimethyl-1,3-benzodioxol-4-yl
- methylcarbamate may be safely used in spot and/or crack and crevice treatments in animal feed handling establishments, including feed manufacturing and processing establishments, such as stores, supermarkets, dairies, meat slaughtering and packing plants, and canneries until the tolerance expiration/revocation date of April 26, 2005.
- (2) The insecticide 2,2-dimethyl-1,3-benzodioxol-4-yl methylcarbamate may be safely used in spot and/or crack and crevice treatments in food handling establishments, including food service, manufacturing and processing establishments, such as restaurants, cafeterias, supermarkets, bakeries, breweries, dairies, meat slaughtering and packing plants, and canneries until the